

WHAT IS CLAIMED IS:

1. A method for sterilizing a biological material that is sensitive to radiation, said method comprising irradiating said biological material with radiation for a time effective to sterilize said biological material at a rate effective to sterilize said biological material and to protect said biological material from said radiation.

2. A method for sterilizing a biological material that is sensitive to radiation, said method comprising:

(i) adding to said biological material at least one stabilizer in an amount effective to protect said biological material from said radiation; and

(ii) irradiating said biological material with a suitable radiation at an effective rate for a time effective to sterilize said biological material.

3. A method for sterilizing a biological material that is sensitive to radiation, said method comprising:

(i) reducing the residual solvent content of said biological material to a level effective to protect said biological material from said radiation; and

(ii) irradiating said biological material with a suitable radiation at an effective rate for a time effective to sterilize said biological material.

4. A method for sterilizing a biological material that is sensitive to radiation, said method comprising:

(i) reducing the temperature of said biological material to a level effective to protect said biological material from said radiation; and

(ii) irradiating said biological material with a suitable radiation at an effective rate for a time effective to sterilize said biological material.

5. A method for sterilizing a biological material that is sensitive to radiation, said method comprising:

0925620-081001

(i) applying to said biological material at least one stabilizing process selected from the group consisting of

- (a) reducing the residual solvent content of said biological material,
- (b) reducing the temperature of said biological material, and
- (c) adding at least one stabilizer to said biological material; and

(ii) irradiating said biological material with a suitable radiation at an effective rate for a time effective to sterilize said biological material, wherein said at least one stabilizing process and the rate of irradiation are together effective to protect said biological material from said radiation.

6. A method for sterilizing a biological material that is sensitive to radiation, said method comprising:

(i) applying to said biological material at least two stabilizing processes selected from the group consisting of

- (a) reducing the residual solvent content of said biological material,
- (b) reducing the temperature of said biological material, and
- (c) adding at least one stabilizer to said biological material; and

(ii) irradiating said biological material with a suitable radiation at an effective rate for a time effective to sterilize said biological material, wherein said at least two stabilizing processes are together effective to protect said biological material from said radiation and further wherein said at least two stabilizing processes may be performed in any order.

7. The method according to claim 3, 5 or 6, wherein said solvent is water.

8. The method according to claim 7, wherein said residual water content is reduced by the addition of an organic solvent.

09925620-081001

9. The method according to claim 3, 5 or 6, wherein said solvent is an organic solvent.
10. The method according to claim 3, 5 or 6, wherein said biological material is suspended in an organic solvent following reduction of said residual solvent content.
11. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said effective rate is not more than about 3.0 kGy/hour.
12. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said effective rate is not more than about 2.0 kGy/hr.
13. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said effective rate is not more than about 1.0 kGy/hr.
14. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said effective rate is not more than about 0.3 kGy/hr.
15. The method according to claim 2, 3, 4, 5 or 6, wherein said effective rate is more than about 3.0 kGy/hour.
16. The method according to claim 2, 3, 4, 5 or 6, wherein said effective rate is at least about 6.0 kGy/hour.
17. The method according to claim 2, 3, 4, 5 or 6, wherein said effective rate is at least about 18.0 kGy/hour.
18. The method according to claim 2, 3, 4, 5 or 6, wherein said effective rate is at least about 30.0 kGy/hour.
19. The method according to claim 2, 3, 4, 5 or 6, wherein said effective rate is at least about 45 kGy/hour.
20. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said biological material is maintained in a low oxygen atmosphere.

0925620-001001

21. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said biological material is maintained in an atmosphere comprising at least one noble gas.

22. The method according to claim 21, wherein said noble gas is argon.

23. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said biological material is maintained in a vacuum.

24. The method according to claim 3, 5 or 6, wherein said residual solvent content is reduced by a method selected from the group consisting of lyophilization, drying, concentration, addition of solute, evaporation, chemical extraction, spray-drying, and vitrification.

25. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 15%.

26. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 10%.

27. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 3%.

28. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 2%.

29. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 1%.

30. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 0.5%.

31. The method according to claim 3, 5 or 6, wherein said residual solvent content is less than about 0.08%.

32. The method according to claim 1, 2, 3, 4, 5 or 6, wherein at least one sensitizer is added to said biological material prior to said step of irradiating said biological material.

09925520-031001

33. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said biological material contains at least one other biological contaminant or pathogen selected from the group consisting of viruses, bacteria and fungi.

34. The method according to claim 2, 5 or 6, wherein said at least one stabilizer is an antioxidant.

35. The method according to claim 2, 5 or 6, wherein said at least one stabilizer is a free radical scavenger.

36. The method according to claim 2, 5 or 6, wherein said at least one stabilizer is a combination stabilizer.

37. The method according to claim 2, 5 or 6, wherein said at least one stabilizer is a ligand.

38. The method according to claim 37, wherein said ligand is heparin.

39. The method according to claim 2, 5 or 6, wherein said at least one stabilizer reduces damage due to reactive oxygen species.

40. The method according to claim 2, 5 or 6, wherein said at least one stabilizer is selected from the group consisting of: ascorbic acid or a salt or ester thereof; glutathione; 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid; uric acid or a salt or ester thereof; methionine; histidine; N-acetyl cysteine; lipoic acid; sodium formaldehyde sulfoxylate; gallic acid or a derivative thereof; propyl gallate and mixtures of two or more thereof.

41. The method according to claim 40, wherein said mixtures of two or more additional stabilizers are selected from the group consisting of: mixtures of ascorbic acid, or a salt or ester thereof, and uric acid, or a salt or ester thereof; mixtures of ascorbic acid, or a salt or ester thereof, and 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid; mixtures of ascorbic acid, or a salt or ester thereof, uric acid, or a salt or ester thereof, and 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid; and mixtures of uric acid, or a salt or ester thereof; lipoic acid; sodium formaldehyde sulfoxylate; gallic acid or a derivative thereof; propyl gallate and 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid.

05925620.001001

42. The method according to claim 2, 5 or 6, wherein said at least one stabilizer is a dipeptide stabilizer.

43. The method according to claim 42, wherein said dipeptide stabilizer is selected from the group consisting of glycyl-glycine (Gly-Gly), carnosine and anserine.

44. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is corpuscular radiation or electromagnetic radiation, or a mixture thereof.

45. The method according to claim 44, wherein said electromagnetic radiation is selected from the group consisting of radio waves, microwaves, visible and invisible light, ultraviolet light, x-ray radiation, gamma radiation and combinations thereof.

46. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is gamma radiation.

47. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is e-beam radiation.

48. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is visible light.

49. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is ultraviolet light.

50. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is x-ray radiation.

51. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is polychromatic visible light.

52. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is infrared.

53. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said radiation is a combination of one or more wavelengths of visible and ultraviolet light.

54. The method according to claim 1, 2, 3, 5 or 6, wherein said irradiation is conducted at ambient temperature.

099255620-081001

55. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said irradiation is conducted at a temperature below ambient temperature.
56. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said irradiation is conducted below the freezing point of said biological material.
57. The method according to claim 1, 2, 3, 4, 5 or 6, wherein said irradiation is conducted below the eutectic point of said biological material.
58. The method according to claim 1, 2, 3, 5 or 6, wherein said irradiation is conducted at a temperature above ambient temperature.
59. A composition comprising at least one biological material and at least one stabilizer in an amount effective to preserve said biological material for its intended use following sterilization with radiation.
60. A composition comprising at least one biological material, wherein the residual solvent content of said biological material is at a level effective to preserve said biological material for its intended use following sterilization with radiation.
61. The composition of claim 60, wherein said residual solvent content is less than about 15%.
62. The composition of claim 60, wherein said residual solvent content is less than about 10%.
63. The composition of claim 60, wherein said residual solvent content is less than about 5%.
64. The composition of claim 60, wherein said residual solvent content is less than about 2%.
65. The composition of claim 60, wherein said residual solvent content is less than about 1%.
66. The composition of claim 60, wherein said residual solvent content is less than about 0.5%.

0925670-081004

67. The composition of claim 60, wherein said residual solvent content is less than about 0.08%.

68. The composition of claim 59 or 60, wherein said biological material is glassy or vitrified.

69. The composition of claim 59 or 60, wherein said biological material is selected from the group consisting of monoclonal immunoglobulins, polyclonal immunoglobulins, glycosidases, sulfatases, urokinase and Factor VIII.

70. The composition of claim 60, wherein the concentration of said biological material is at least about 0.5%.

71. The composition of claim 60, wherein the concentration of said biological material is at least about 1%.

72. The composition of claim 60, wherein the concentration of said biological material is at least about 5%.

73. The composition of claim 60, wherein the concentration of said biological material is at least about 10%.

74. The composition of claim 60, wherein the concentration of said biological material is at least about 15%.

75. The composition of claim 60, wherein the concentration of said biological material is at least about 20%.

76. The composition of claim 60, wherein the concentration of said biological material is at least about 25%.

77. The composition of claim 60, wherein the concentration of said biological material is at least about 50%.

78. The method according to claim 32, wherein said sensitizer is a ligand.

79. The method according to claim 78, wherein said ligand is a metal ion.

80. The method according to claim 79, wherein said metal ion is a copper ion.

099255620-08-1001